

FOR TEMPORARY EXTERNAL USE





Antibacterial Burn Wrap

5in x 5ft Dressing



For the Pre-Hospital Management of 1st and 2nd Degree Burns with Aquanova Ag Technology

Each dressing contains nominally 0.75% by weight of ionic silver

INSTRUCTIONS FOR USE ON BACK



















For the Pre-Hospital Management of 1st and 2nd Degree Burns

Instructions for Use:

- 1. Irrigate wound area according to locally established protocols.
- 2. Tear open dressing pack. Unravel Silvapro Antibacterial Burn Wrap and cover burn, overlapping at least 2 inches onto the surrounding skin.
- 3. For exuding wounds Silvapro Antibacterial Burn Wrap can be applied dry and gels on contact with wound fluid to form an antibacterial barrier, or alternatively, the dressing should be wetted after application using saline or water to cool the burn.
- 4. Cover Silvapro Antibacterial Burn Wrap with a sterile loosely woven gauze and secure.
- 5. Remove the gauze cover dressing periodically and inspect the Silvapro Antibacterial Burn Wrap while it remains in place on the burn.

Once product has been used:

Seek MEDICAL CARE as soon as possible. Show empty pack to medical personnel. Discard unused portion of the Silvapro Antibacterial Burn Wrap after dressing the wound.

Contraindications: The dressing is contraindicated for third (3rd) degree burns or for use on individuals with a known sensitivity to silver or chitosan.

Warnings: Device is not intended for Implantation. For temporary external use only. Do not apply over eyes. Frequent or prolonged use of this product may result in permanent discoloration of skin, due to the ionic silver. Contains chitosan from shellfish. Allergy studies show no adverse reaction to date. Data on file at Medtrade Products Ltd.

SILVAPRO™ is a trademark of Medtrade Products Ltd. All rights reserved.

In-vitro antibacterial data has shown that Silvapro Escherichia coli, Pseudomonas aeruginosa, Candi albicans, Enterococcus faecalis (VRE),

> 50 x 20 mm UDI 2D barcode and data