Antibacterial Burn Dressing

5in x 9in Dressing



Dressina covers approx. 2% total body surface area

For the Pre-Hospital Management of 1st and 2nd Degree Burns with Aguanova Ag Technology

Each dressing contains nominally 0.75% by weight of ionic Silver

INSTRUCTIONS FOR USE ON BACK















For The Pre-Hospital Management of 1st and 2nd Degree Burns



Instructions for Use:

- 1. Irrigate wound area according to locally established protocols.
- 2. Tear open dressing pack. Unravel Silvapro Antibacterial Burn Dressing and cover burn. Overlapping at least 2inches onto the surrounding skin.
- For exuding wounds Silvapro Antibacterial Burn Dressing can be applied dry and gels on contact with wound fluid to form an antibacterial barrier, or alternatively, the dressing should be wetted after application using saline or water to cool the burn.
 Cover Silvapro Antibacterial Burn Dressing with a sterile loosely woven gauze and

Once product has been used:

Seek MEDICAL CARE as soon as possible. Show empty pack to medical personnel. Discard unused portion of the Silvapro Antibacterial Burn Dressing after dressing the wound.

Contraindication:

The dressing is contraindicated for third (3rd) degree burns or for use on individuals with known sensitivity to ionic silver.

external use only. Do not apply over eyes. Frequent or prolonged use of this product may result in permanent discoloration of skin, due to the ionic silver. Contains chitosan from shellfish. Allergy studies show no adverse reaction to date. D<u>ata on file at</u> Medtrade Products Ltd



Manufactured by: Medtrade Products Ltd.

SILVAPRO™ is a trademark of Medtrade Products Ltd. All rights reserved.



In-vitro antibacterial data has shown that Silvapro

50 x 20 mm UDI 2D barcode and data



REF FG0882519









